

# The free and bound forms of Lpp occupy distinct subcellular locations in *Escherichia coli*

Charles E. Cowles,<sup>1</sup> Yongfeng Li,<sup>2†</sup>  
Martin F. Semmelhack,<sup>2</sup> Ileana M. Cristea<sup>1</sup> and  
Thomas J. Silhavy<sup>1\*</sup>

Departments of <sup>1</sup>Molecular Biology and <sup>2</sup>Chemistry,  
Princeton University, Princeton, NJ 08544, USA.

## Summary

The lipoprotein Lpp is the most numerically abundant protein in *Escherichia coli*, has been investigated for over 40 years, and has served as the paradigmatic bacterial lipoprotein since its initial discovery. It exists in two distinct forms: a ‘bound-form’, which is covalently bound to the cell’s peptidoglycan layer, and a ‘free-form’, which is not. Although it is known that the carboxyl-terminus of bound-form Lpp is located in the periplasm, the precise location of free-form Lpp has never been determined. For decades, it has been widely assumed that free-form Lpp is associated with bound-form. In this work, we show that the free and bound forms of Lpp are not largely associated with each other, but are found in distinct subcellular locations. Our results indicate that free-form Lpp spans the outer membrane and is surface-exposed, whereas bound-form Lpp resides in the periplasm. Thus, Lpp represents a novel example of a single lipoprotein that is able to occupy distinct subcellular locations, and challenges models in which the free and bound forms of Lpp are assumed to be associated with each other.

## Introduction


Surface-exposed lipoproteins play important biological roles in many Gram-negative pathogens (Cornelissen *et al.*, 1998; Dashper *et al.*, 2000; Jin *et al.*, 2001; Leuzzi *et al.*, 2005). Even though three lipoproteins have been demonstrated to be surface-exposed in *Escherichia coli* (Manning *et al.*, 1980; Drummelsmith and Whitfield, 2000; Robinson *et al.*, 2006), it is commonly assumed in many descriptions of this model organism’s cell envelope that

the vast majority of lipoproteins reside in the periplasm (Ruiz *et al.*, 2006; Tokuda, 2009). Surface proteolysis of intact cells has long been the favoured method to demonstrate a protein’s surface-exposure, but *E. coli* has evolved to thrive in the protease-rich environment of the mammalian intestinal tract. Indeed, many surface-exposed proteins in *E. coli* are protease-resistant in intact cells (Behr *et al.*, 1980; Manning *et al.*, 1980; Cole *et al.*, 1983; Ronco *et al.*, 1988; Bolla *et al.*, 1990). Moreover, none of the surface-exposed lipoproteins in *E. coli* was discovered using protease-based methods (Manning *et al.*, 1980; Drummelsmith and Whitfield, 2000; Robinson *et al.*, 2006). Thus, novel, protease-independent methods must be developed in order to adequately test the prevalence of surface-exposed lipoproteins in *E. coli*.

Lpp (a.k.a. lipoprotein, Braun’s lipoprotein, murein lipoprotein and MlpA) is the best-characterized bacterial lipoprotein. It is a small, outer membrane (OM) lipoprotein in *E. coli* that physically tethers the OM to the peptidoglycan layer (Braun and Rehn, 1969; Braun and Sieglin, 1970; Braun and Wolff, 1970; Braun and Bosch, 1972). Lpp is estimated to be present at as many as 750 000 copies per cell, making it the most numerically abundant protein in *E. coli* (Neidhardt, 1996; Nikaido, 1996). Cells lacking Lpp exhibit numerous OM defects, including increased OM permeability to antibiotics and other toxic small molecules, leakage of periplasmic contents into the extracellular environment, and heightened production of outer membrane vesicles (Hirota *et al.*, 1977; Yem and Wu, 1977; Suzuki *et al.*, 1978).

*In vivo*, Lpp exists in two separate forms: a ‘bound-form’, which is covalently attached by its carboxyl-terminal lysine to the peptidoglycan layer, and a ‘free-form’, which is not attached (Braun and Rehn, 1969; Braun and Bosch, 1972; Inouye *et al.*, 1972). In the cell, the free and bound forms of Lpp exist in an approximate 2:1 ratio respectively (Inouye *et al.*, 1972). Although it has long been known that the carboxyl-terminus of bound-form Lpp is located in the periplasm, published models omit description of the location of free-form Lpp (Nikaido and Nakae, 1979; Nikaido, 1996), or predict it to be physically associated with the bound form of the protein (Inouye, 1974; Shu *et al.*, 2000). However, a physical association *in vivo* between free-form and bound-form Lpp subunits has never been demonstrated, and the location of free-form Lpp remains unclear.

Accepted 2 January, 2011. \*For correspondence. E-mail tsilhavy@princeton.edu; Tel. (+1) 609 258 5899; Fax (+1) 609 258 2957.  
†Present address: Rm. 1306 #11, Lon 97, Hindan Road, Shanghai, China.



**Fig. 1.** A method for labelling surface-exposed proteins in *E. coli*.

A. Structure of NHS-LC-LC-biotin.

B. Western blot probed for biotin of soluble (sol.) and insoluble (insol.) fraction proteins from whole cells labelled with NHS-LC-LC-biotin.

C. Western blot probed for biotin of soluble (sol.) and insoluble (insol.) fraction proteins from a lysate labelled with NHS-LC-LC-biotin.

D. Western blot probed for biotin of cellular proteins from wild-type, 'wt'; *ompA*-null, '*ompA*'; and *lpp*-null, '*lpp*' strains. The protein band that is suspected to be biotin-labelled OmpA is indicated with an asterisk.

Here, we describe a widely applicable method to label surface-exposed proteins in *E. coli*. Surprisingly, our results show that free-form Lpp is surface-exposed. We provide evidence that bound- and free-form Lpp occupy two separate and distinct subcellular locations: bound-form Lpp is located entirely within the periplasm, whereas free-form Lpp represents a transmembrane topology that spans the OM and is exposed at the cell surface. As such, Lpp provides a unique example of a lipoprotein that stably occupies two distinct subcellular locations: one, which spans the OM lipid bilayer, and another, which exists in the periplasm.

## Results

### *A method for labelling surface-exposed proteins in E. coli*

Numerous surface-exposed *E. coli* OM proteins are highly resistant to proteolysis from the extracellular surface

(Behr *et al.*, 1980; Manning *et al.*, 1980; Cole *et al.*, 1983; Ronco *et al.*, 1988; Bolla *et al.*, 1990). An alternative method to identify surface-exposed proteins lies in labelling cells with OM-impermeable reagents; however, this strategy requires careful consideration of the unique barrier properties of the OM. In general, the Gram-negative OM functions as a barrier to prevent the entry of hydrophobic compounds of any molecular weight, as well as high-molecular-weight, hydrophilic compounds, while also allowing the non-specific entry of low-molecular-weight, hydrophilic compounds (Nikaido and Rosenberg, 1981; Nikaido, 2003).

We sought to find a protein-labelling reagent that would not cross the OM and enter the periplasm. NHS-LC-LC-biotin is a hydrophobic, primary amine-reactive biotinylation reagent (Fig. 1A). Although the manufacturer describes this reagent as being generically membrane-permeable, we reasoned that the *E. coli* OM might

exclude it, owing to its hydrophobicity and relatively high molecular weight. If this reagent does enter the periplasm, then it should not be able to label OM proteins with a periplasmic orientation, without also labelling some of the more abundant soluble periplasmic proteins.

To examine the OM permeability of NHS-LC-LC-biotin, we labelled whole cells and examined the distribution of the biotin label in samples that were subsequently fractionated into soluble (containing mostly cytoplasmic and periplasmic proteins) and insoluble (containing mostly membrane proteins) fractions. Because the label was almost exclusively detected in insoluble-fraction proteins (Fig. 1B), we concluded that the labelling reagent may not have entered the periplasm to a significant degree.

We were concerned that this reagent might selectively partition into membrane lipid bilayers or selectively label membrane proteins for unforeseen reasons, two situations that could also yield the observed result (Fig. 1B). To test these possibilities, we labelled a cell-free lysate and subsequently separated the lysate into the soluble and insoluble fractions, as above. Upon doing so, we observed numerous, novel labelled proteins in the soluble fraction, thereby eliminating the possibilities that NHS-LC-LC-biotin selectively partitions into membranes or labels only insoluble proteins (Fig. 1C).

Even though our controls seemed to indicate that NHS-LC-LC-biotin labels only surface-exposed proteins, we were deeply concerned that our labelling method could damage or disrupt the OM permeability barrier. To examine the effects of our labelling method on OM permeability, we compared labelled and unlabelled cells in several ways. However, labelled cells showed no detectable decrease in viability (as measured by determining the efficiency of plating on solid SDS/EDTA medium) or growth rate, as compared with unlabelled controls (data not shown). In addition, labelled and unlabelled cells had indistinguishable zones of clearing around EDTA-treated filter discs placed in the centre of solid media containing SDS (data not shown). Finally, cells labelled by our labelling method were no more permeable to Sytox Green than their unlabelled counterparts (77/2289 labelled cells were permeable to Sytox Green, as compared with 101/2048 unlabelled cells). These results demonstrate that our labelling method does not damage or disrupt the *E. coli* OM permeability barrier in a detectable way.

To further investigate the surface specificity of NHS-LC-LC-biotin, we sought to examine known surface-exposed proteins using null mutants lacking these proteins. One such protein is OmpA, an abundant transmembrane  $\beta$ -barrel protein that serves as a surface receptor for Tull\* bacteriophage (Datta *et al.*, 1977). By comparing the biotin label profiles of wild-type and *ompA*-null strains, we observed the loss of a major surface-exposed protein in the *ompA*-null strain that migrates with the same electro-

phoretic mobility as OmpA (Fig. 1D). We were also interested in the identity of the low-molecular-weight protein that is intensely labelled by this reagent. Lpp is an extremely abundant, low-molecular-weight OM protein with this same electrophoretic mobility. Indeed, an *lpp*-null strain showed no labelled proteins with this electrophoretic mobility (Fig. 1D). These results suggest that at least a portion of the Lpp molecules in *E. coli* are surface-exposed.


#### *A soluble, high-molecular-weight labelling reagent also labels Lpp*

Lpp was not previously thought to be surface-exposed in *E. coli*. Therefore, although our control experiments indicate that NHS-LC-LC-biotin labels only surface-exposed molecules, and that Lpp is labelled by this reagent, we were hesitant to conclude that Lpp is truly surface-exposed. To more rigorously test whether Lpp is surface-exposed, we synthesized a novel, primary-amine-reactive, hydrophilic, high-molecular-weight labelling reagent, PEG-NHS-biotin. The polyethyleneglycol backbone of this molecule makes it hydrophilic, yet of sufficiently high molecular weight so as to preclude its entry into the periplasm (Nakae and Nikaido, 1975; Decad and Nikaido, 1976) (Fig. 2A). After labelling intact cells with PEG-NHS-biotin and fractionating the sample into soluble and insoluble fractions, we again observed the vast majority of the label in insoluble fraction proteins with a pattern similar to the pattern of proteins labelled by NHS-LC-LC-biotin (Fig. 2B). Moreover, Lpp was still intensely labelled by this reagent.

Because PEG-NHS-biotin labels proteins less intensely than NHS-LC-LC-biotin, we used longer exposure times in developing our blots. In doing so, we observed a single biotinylated protein in the soluble fraction. However, this protein was found to share a similar electrophoretic mobility to that of BCCP, a cytoplasmic biotin carrier protein with a covalently attached biotinyl prosthetic group (Li and Cronan, 1992). Furthermore, this protein was observed to be uniformly biotinylated in unlabelled cells (data not shown). Thus, we can conclude that PEG-NHS-biotin cannot enter the *E. coli* periplasm. Because two different primary amine-reactive labelling reagents that we predicted to be OM-impermeable for different reasons both labelled Lpp, we more seriously considered the possibility that some of the Lpp molecules in *E. coli* may, in fact, be surface-exposed.

#### *The LC-LC-biotin label can be removed by surface proteolysis*

The LC-LC-biotin moiety has two internal isopeptide bonds and creates a third when it attaches to a primary



**Fig. 2.** Lpp is surface-exposed in *E. coli*.  
**A.** Structure of PEG-NHS-biotin. An identical *N*-hydroxysuccinimidyl-biotin moiety is found at either end of the PEG 3500 backbone.  
**B.** Western blot probed for biotin of soluble and insoluble fraction proteins from whole cells labelled with PEG-NHS-biotin.  
**C.** Western blots, cropped to show the OmpA and Lpp regions, probed for specific proteins or biotin, as indicated. Samples were frozen immediately after labelling ( $T_0$ ), or treated with no protease ( $\emptyset$ ), proteinase K (PK) or trypsin (Tr), as indicated.  
**D.** Western blot probed for biotin of cells expressing the indicated quantities (in ng) of NlpB or TraT, labelled with NHS-LC-LC-biotin. The known positions of NlpB and TraT are indicated to the right.


amine. Although *E. coli* surface-exposed proteins are highly resistant to proteolysis by extracellular proteases, we postulated that the surface-exposed LC-LC-biotinyl group might be protease-sensitive. Indeed, when we treated intact LC-LC-biotinylated cells with either proteinase K or trypsin, significant amounts of the biotin label were removed from both OmpA and Lpp (Fig. 2C) with no concomitant proteolytic degradation of either of the proteins to which it was attached. As a control, OmpA is especially informative in that it is known to be susceptible to proteolysis from the periplasmic face of the OM but not from the extracellular surface (Chen *et al.*, 1980). Since OmpA levels were not affected by the protease treatment, we can eliminate the possibilities that the proteases entered the periplasmic space or that the LC-LC-biotin modification renders proteins susceptible to surface proteolysis. Therefore, we conclude that NHS-LC-LC-biotin primarily labels surface-exposed molecules and, additionally, that a portion of the Lpp molecules in *E. coli* are surface-exposed.

#### *A labelling control using lipoproteins with known topologies*

To determine if NHS-LC-LC-biotin can penetrate the OM and label periplasmic lipoproteins from this location, we examined whether we could label a known periplasmic lipoprotein, NlpB (Wu *et al.*, 2005), or a known surface-exposed lipoprotein, TraT (Manning *et al.*, 1980). Although TraT was clearly labelled, we were unable to detect labelled NlpB, even when this protein was produced at over sixfold higher levels (Fig. 2D). These results establish that NHS-LC-LC-biotin specifically labels surface-exposed lipoproteins.

#### *A label-independent confirmation of Lpp surface-exposure*

In other organisms, surface-exposed lipoproteins can deliver fluorescent proteins to the cell surface (Schulze and Zuckert, 2006). If Lpp can function similarly to deliver an epitope tag to the cell surface, it would provide a



**Fig. 3.** Label-independent confirmation of Lpp surface-exposure. **A.** Western blot of Lpp from cells expressing Lpp-FLAG, separated into the free-form (FF) and bound-form (BF) subcellular fractions. The FF sample was diluted 10-fold prior to loading. **B.** Western blots of whole cells expressing native Lpp or Lpp-FLAG untreated (-) or treated (+) with trypsin, as indicated. Blots were probed for OmpA, Lpp or the FLAG epitope, as indicated. The novel Lpp species that results from trypsin treatment of the Lpp-FLAG protein is indicated with an asterisk. **C.** Western blot probed for biotin of whole cells expressing native Lpp or Lpp-FLAG untreated (-) or treated (+) with trypsin, as indicated, then labelled with biotin. The novel Lpp species that results from trypsin treatment of the Lpp-FLAG protein is indicated with an asterisk.

labelling reagent-independent means to confirm the surface-exposure of free-form Lpp. Even though the Lpp-FLAG construct has a non-native carboxyl-terminus (DYKDDDDK), the epitope-tagged protein was present in cells as both the free and bound forms (Fig. 3A). Although

the amount of bound-form is quite low (bound-form Lpp migrates more slowly than free-form, usually as a ladder of bands, because a heterogeneous population of peptidoglycan fragments are attached to its carboxyl-terminus), this allele complemented an *lpp*-null strain for SDS/EDTA resistance on solid media as well as the wild-type allele when expressed from a plasmid, indicating that it is physiologically functional (data not shown).


Even though Lpp is not susceptible to proteolysis from the extracellular surface (Fig. 2C), we discovered that we could remove the FLAG epitope from Lpp-FLAG using whole cell surface proteolysis. Indeed, we were able to use trypsin to remove the epitope from approximately one-third of the cells' Lpp-FLAG molecules without affecting OmpA (Fig. 3B). We conclude from these results that the carboxyl-terminus of free-form Lpp is exposed at the cell surface.

We wished to determine whether our inability to remove the FLAG epitope from all of the cells' Lpp-FLAG molecules was due to incomplete proteolysis, the FLAG epitope interfering with the protein's localization to the cell surface, or a reflection of the native distribution of the protein in the cell. To address this, we labelled intact, surface-proteolysed Lpp-FLAG-expressing cells with NHS-LC-LC-biotin to determine whether both the epitope-containing and truncated forms of the protein could be labelled. Both forms of the protein were labelled by LC-LC-biotin (Fig. 3C), thereby suggesting that our inability to remove the FLAG epitope from all of the cells' Lpp-FLAG proteins is simply due to incomplete proteolysis.

#### *The free and bound forms of Lpp exist in different subcellular compartments*

Lpp exists in two different forms in *E. coli*: a 'bound-form' in which the protein's carboxyl-terminus is covalently attached to the peptidoglycan layer, and a 'free-form', which is not (Inouye *et al.*, 1972). Because bound-form Lpp is known to be located in the periplasm, surface-labelling reagents should not label it. To test this, we labelled cells with NHS-LC-LC-biotin, fractionated the two forms of Lpp, and compared the amount of biotin label attached to each of the two forms, relative to the amount of Lpp protein present in each fraction. We found that the biotin label was predominantly associated with free-form Lpp (Fig. 4A). This result provides further evidence that NHS-LC-LC-biotin cannot readily enter the *E. coli* periplasm.

We considered the possibility that bound-form Lpp might not be susceptible to labelling by NHS-LC-LC-biotin *in vivo*. To test this, we labelled cells with a high concentration of NHS-LC-LC-biotin and overexposed blots to determine whether we could detect any biotin-labelled



**Fig. 4.** The free and bound forms of Lpp occupy distinct subcellular locations.  
 A. Western blots of samples from cells labelled with NHS-LC-LC-biotin and separated into free-form (FF) and bound-form (BF) Lpp fractions. Blots were probed for Lpp or biotin, as indicated.  
 B. Western blots of samples from wild-type (wt) or *imp4213* (*imp*) cells labelled with a high concentration of NHS-LC-LC-biotin and separated into free-form (FF) and bound-form (BF) Lpp fractions. Blots were probed for Lpp or biotin, as indicated.  
 C. Western blots of Lpp from EDTA-treated cells left untreated (-) or treated (+) with proteinase K (PK) and separated into free- and bound-form Lpp fractions.

bound-form Lpp, whatsoever. Indeed, under these conditions we observed biotin-labelled bound-form Lpp, thereby indicating that bound-form Lpp can be modified by NHS-LC-LC-biotin *in vivo* (Fig. 4B). Furthermore, bound-form Lpp prepared from a similarly labelled *imp4213* strain (a strain with severe OM permeability defects) showed noticeably more biotin label than did bound-form from the wild-type strain (Fig. 4B) (Braun and


Silhavy, 2002). These results provide additional evidence that NHS-LC-LC-biotin is excluded by the OM in wild-type strains. More importantly, these results suggest that the label's bias towards the free-form is due to a topological difference between the two forms.

Our results indicate that at least some of the cell's free-form Lpp molecules are labelled by OM-impermeable labelling compounds (Fig. 4A), yet we also observed that free-form Lpp is insensitive to proteolysis from the extracellular surface (Fig. 2C). In contrast, bound-form Lpp is not labelled by OM-impermeable labelling compounds to a significant degree. Therefore, bound-form Lpp, which is periplasmic, as well as any periplasmic free-form Lpp molecules, should be sensitive to proteases introduced into the periplasm. To test this possibility, we disrupted cells' OM permeability barrier with EDTA and passed proteinase K into the periplasm. Upon doing so, we observed degradation of only bound-form Lpp, whereas free-form Lpp was not noticeably affected (Fig. 4C). Thus, both label-dependent and label-independent methods indicate that the free and bound forms of Lpp occupy distinct subcellular compartments.

#### *The carboxyl-terminus of Lpp is surface-exposed*

In order to better understand the orientation and position of the free-form Lpp molecules that are labelled by our method, we employed mass spectrometry (MS) to identify where LC-LC-biotin attaches to the protein. Three LC-LC-biotin-modified peptides, LDNMATK\*, K\* and LDNMATK\*YRK\* (where '\*' denotes modification by LC-LC-biotin), were detected by MALDI LTQ Orbitrap MS analysis with accuracies of 0.8, 0 and 1.3 p.p.m. respectively. Thus, the two carboxyl-terminal lysines of Lpp (Lys<sub>55</sub> and Lys<sub>58</sub>) seem to be modified with LC-LC-biotin (Fig. 5A). To verify that Lpp is modified at these positions, we subjected the dually LC-LC-biotin-modified tryptic peptide corresponding to the 10, carboxyl-terminal amino acid residues (LDNMATK\*YRK\*, Fig. 5A) to MS<sup>2</sup> and MS<sup>3</sup> analyses. The resulting fragment ions confirmed the presence of LC-LC-biotin modifications attached to the two carboxyl-terminal lysines (Fig. S1).

Our MS analyses indicate that the carboxyl-terminus of free-form Lpp is exposed at the cell surface, and suggest that most of the remainder of the protein is not. However, we were unable to detect all of the tryptic peptides by this method, including the acylated, amino-terminal tryptic lipopeptide. Because the amino-terminal tryptic lipopeptide ends at a lysine residue, it is a potentially modifiable position of the protein. Thus, our MALDI MS analysis results alone cannot exclude the possibility that Lpp also exists in an orientation with its amino-terminus exposed at the cell surface. In addition, it is possible that other positions of the protein are labelled, but not observed in our



**Fig. 5.** The carboxy-terminus of Lpp is surface exposed.

A. Schematic of Lpp depicting the acylated amino-terminus, internal deletion from the Lpp $_{\Delta 17-37}$  mutant (central grey box), deleted carboxy-terminal lysine from the Lpp $_{\Delta K58}$  mutant (boxed in grey at right), and peptide that was isolated and used in MS<sup>2</sup> and MS<sup>3</sup> analyses (underlined). Asterisks indicate the lysine residues modified by LC-LC-biotin. B. Western blots of LC-LC-biotin-labelled *lpp*-null cells expressing different forms of Lpp. Blots were probed for Lpp or biotin, as indicated. Cells harboured pBAD18 plasmid vector (v), pBAD/*lpp* (wt), pBAD/*lpp* $_{\Delta K58}$  ( $\Delta K58$ ) or pBAD/*lpp* $_{\Delta 55-58}$  ( $\Delta 55-58$ ).

MALDI MS experiment due to low levels or other unforeseen reasons.

To further investigate whether Lys<sub>55</sub> and Lys<sub>58</sub> are the only labelled positions of the protein, we cloned and expressed wild-type *lpp*, an *lpp* allele lacking the carboxyl-terminal lysine (*lpp* $_{\Delta K58}$ ), and an *lpp* allele lacking the four carboxyl-terminal-most amino acids, KYRK (*lpp* $_{\Delta 55-58}$ ). When expressed at equal levels in an *lpp*-null strain (Fig. 5B), wild-type Lpp showed the highest level of biotin labelling, whereas the variant lacking the carboxyl-terminal-most lysine showed reduced levels of labelling (Fig. 5B). In agreement with our MALDI MS/MS results, the Lpp variant that lacks the four carboxyl-terminal-most amino acids showed no detectable biotin labelling (Fig. 5B). From these results, we conclude that surface-exposed free-form Lpp molecules adopt a topology with their carboxyl-termini, but not their amino-termini, exposed at the cell surface.

#### Mutant *lpp* alleles alter the levels of surface-exposed Lpp

Two different mutant alleles of *lpp* that produce either entirely free-form or mostly bound-form Lpp have been described (Zhang *et al.*, 1992; Zhang and Wu, 1992). By reconstructing these alleles in an isogenic strain background at the native chromosomal locus, free- and bound-form Lpp levels can be altered to examine the effects on its surface-exposure.

Bound-form Lpp is covalently bound to the cell's peptidoglycan layer via its carboxyl-terminal lysine residue by a linkage between its  $\epsilon$ -amino group and the carboxyl group at the L-centre of meso-diaminopimelate (Braun and Wolff, 1970). Thus, the mutant allele lacking this lysine codon produces a protein that exists entirely as the free


form (Zhang and Wu, 1992). As expected, a strain carrying this allele (*lpp* $_{\Delta K58}$ ) expressed only free-form Lpp (Fig. 6A).

The *lpp* allele that expresses mostly bound-form Lpp has an internal deletion of codons 17–37 of the mature, wild-type protein (Fig. 5A). In agreement with a previous study (Zhang *et al.*, 1992), a strain carrying this allele (*lpp* $_{\Delta 17-37}$ ) produced almost entirely bound-form Lpp with very little free-form (Fig. 6A). To determine whether the levels of free-form Lpp in these strains were reflective of the amount of surface-exposed Lpp, we surface-labelled cells with NHS-LC-LC-biotin and examined the amount of biotinylated Lpp. As expected if only free-form Lpp is surface-exposed, the extent of the biotin labelling directly paralleled the levels of free-form Lpp in each of these strains (Fig. 6B).

Lpp levels in the *lpp* $_{\Delta 17-37}$  strain appear lower than in wild-type when examined by immunoblot (Fig. 6A). However, we know from metabolic labelling experiments that the mutant and wild-type proteins are produced at similar levels (See Supplementary Information). The fact that the mutant protein is missing 1/3 of the amino acids that are present in the wild-type protein likely explains the observed difference in antibody binding.

#### The oligomeric state of free-form Lpp

A physical association *in vivo* between free- and bound-form Lpp subunits has not been reported, although a trimer comprised of only free-form Lpp subunits has been observed (Choi *et al.*, 1986). If the free and bound forms of Lpp are physically associated with each other *in vivo*, and if a substantial number of trimers have one or more bound-form Lpp monomers present, then the number of cross-linked trimers will increase substantially if covalent



**Fig. 6.** Mutant alleles alter levels of surface-exposed Lpp.

A. Western blot probed for Lpp of samples from cells expressing the indicated *lpp* alleles, separated into the free-form (FF) and bound-form (BF) subcellular fractions. Because the Lpp $_{\Delta 17-37}$  mutant protein is not detected well by the antibody used in these studies, twice the sample volume was loaded for this strain, relative to the amounts loaded from the other 2 strains.

B. Western blot probed for biotin of samples from cells expressing the indicated forms of Lpp, labelled with LC-LC-biotin. Equal volumes of cell density-normalized samples were loaded in each lane.

C. Western blot of Lpp from cells expressing the indicated forms of Lpp, cross-linked with the indicated concentrations of disuccinimidyl suberate (DSS). The apparent monomeric (m), dimeric (d) and trimeric (t) complexes are indicated.

binding of Lpp to the peptidoglycan is prevented. However, if the two forms are not physically associated, then preventing Lpp from being covalently bound to the peptidoglycan should have little effect on the levels of cross-linked free-form Lpp trimer. It is important to note that in the absence of lysozyme treatment to digest the cell's peptidoglycan layer, bound-form Lpp does not enter an acrylamide gel. Therefore, by not treating samples with lysozyme, one can be certain that only free-form Lpp will be observed. To investigate the potential physical association between free- and bound-form Lpp *in vivo*, we exposed wild-type and *lpp* $_{\Delta K58}$  cells to increasing concentrations of disuccinimidyl suberate and examined Lpp trimer levels by immunoblot analysis. Both strains revealed similar amounts of the free-form trimer that increased as the level of cross-linker was increased, confirming that the free and bound forms of Lpp are not largely associated *in vivo* (Fig. 6C). Slightly more Lpp trimer was observed in the *lpp* $_{\Delta K58}$  strain at all cross-linker concentrations, likely reflecting the fact that this strain produces approximately 50% more Lpp than the wild-type.

## Discussion

It has long been known that there are two forms of Lpp: a 'bound-form,' which is covalently bound to the peptidoglycan, and a 'free-form,' which is not (Braun and Rehn, 1969; Braun and Bosch, 1972; Inouye *et al.*, 1972). Some studies have assumed that the two forms are associated with each other (Inouye, 1974; Shu *et al.*, 2000); however, such an association has never been experimentally demonstrated. Instead, we have discovered that most or all of the free-form molecules are surface exposed. Free-form Lpp can be labelled by an NHS-activated biotinylation reagent, which is excluded by the OM, whereas bound-form labels poorly. Conversely, bound-form Lpp can be degraded by proteases introduced into the periplasm, whereas free-form cannot. If a FLAG tag is added at the carboxyl-terminus of Lpp, the mutant protein still retains its biological function. Most strikingly, the tagged protein is covalently attached to the peptidoglycan, and a substantial fraction of the FLAG-tagged molecules are surface-exposed. The simplest explanation for all of these results is that the free and bound forms of Lpp are not

largely associated with each other, but instead occupy distinct subcellular locations.

When biosynthesis of Lpp is complete, the molecule is tethered to the periplasmic face of the inner membrane by its amino-terminal lipid moieties. The Lol system then transports individual Lpp monomers across the periplasm to the OM, leaving them tethered to the inner surface of the OM (Yokota *et al.*, 1999). Our data indicate that the carboxyl-terminus, but not the amino-terminus of free-form Lpp, is exposed at the cell surface. Accordingly, we suspect that the lipid moieties remain embedded in the inner leaflet of the OM and that the molecule assumes a transmembrane conformation. Such a topology would explain why free-form Lpp is resistant to proteases present on either side of the OM. Because Lpp is present in two distinct subcellular locations, we suggest that it may be more accurate to refer to the 'periplasmic' and 'transmembrane' forms of Lpp in certain contexts.

Although it may be very transient, all bound-form molecules must initially exist as periplasmic free-form molecules, prior to being covalently bound to the peptidoglycan layer. In addition, it is possible that a small fraction of the cell's free-form Lpp molecules remain in the periplasm, perhaps associated with bound-form molecules. In either case, it is possible that a small subpopulation of the cell's free-form molecules resides in the periplasm. It is important to note that we do not suspect that we would be able to detect this putative subpopulation with the experiments that we have performed. Nevertheless, our results suggest that the vast majority of the free-form molecules in *E. coli* are exposed at the cell surface. On average, *E. coli* cells express as many as 500 000 free-form Lpp monomers. Considered from the perspective that there are approximately 1.4–3.5 million copies of LPS per cell in the OM (Neidhardt, 1996; Nikaido, 1996), our results imply that free-form Lpp molecules are a significant feature of the *E. coli* OM bilayer.

We do not yet understand how free-form Lpp is inserted into the OM. It is possible that the Bam complex, which assembles  $\beta$ -barrel proteins in the OM, is involved (Ruiz *et al.*, 2006). Alternatively, there may be a dedicated translocase or a folding catalyst that can interconvert the periplasmic and transmembrane forms before the former is covalently attached to the cell wall. Given the small size of Lpp it is also possible that membrane insertion happens spontaneously. Regardless of the mechanism, it is clear that it can translocate peptide tags, such as the FLAG epitope (DYKDDDDK), that are attached to the carboxyl-terminus of the molecule. This is particularly striking because the FLAG epitope is highly charged and, thus, would likely present a challenging translocation substrate.

Previous reports indicated that free-form Lpp is trimeric (Choi *et al.*, 1986; Shu *et al.*, 2000). We have confirmed

this result and have extended it to show that the transmembrane form is also likely to be trimeric. We do not know the structure of the transmembrane form. However, the amino acid sequence of Lpp is not incompatible with a transmembrane form. Indeed, models depicting oligomers of the protein in transmembrane form have been proposed (Inouye, 1974). The crystal structure of an unlipidated Lpp peptide, corresponding to residues 2–57 of the mature protein, has been determined and reveals a trimeric, coiled-coil structure (Shu *et al.*, 2000). This trimer's surface is studded with numerous polar and charged residues and it is unlikely that such a structure could stably span the OM. However, it is possible that this structure represents the periplasmic topology of Lpp. Regardless of the structure, our results suggest that most of the periplasmic monomers are cross-linked to the cell wall because we see little evidence indicating substantial levels of periplasmic free-form Lpp molecules. Unfortunately, since the oligomeric state of the periplasmic form of Lpp is not known the biological significance of the published structure is not clear.

Bound-form Lpp tethers the OM to the peptidoglycan cell wall, and pronounced OM defects are apparent in strains carrying the *lpp* $\Delta$ <sub>K58</sub> mutation, which breaks this tether, but leaves free-form undisturbed (Zhang and Wu, 1992). One way to address the function of free-form Lpp would be to phenotypically characterize a strain that produces only bound-form Lpp. As noted above, the *lpp* $\Delta$ <sub>17–37</sub> mutation virtually eliminates free-form Lpp. However, this mutation confers a mucoid phenotype on solid media, and is dominant in diploid analysis, demonstrating that the truncated mutant protein is toxic. Thus, the physiological function of free-form Lpp remains unclear. It may be that cells produce an excess of free-form Lpp simply to ensure sufficient levels of the protein are available to be covalently bound to the peptidoglycan. It is also possible that the transmembrane form of the protein can be converted into the periplasmic form. Therefore, the transmembrane topology may represent a neutral location where reserves of free-form Lpp can be stored for immediate use at a later time.

We do not understand what determines the ratio of the periplasmic and transmembrane forms. There may be a regulatory mechanism to evaluate and adjust this ratio or it may be an intrinsic property of the protein, itself. Our results with the *lpp* $\Delta$ <sub>K58</sub> mutation suggest that the transmembrane form is the default state. This may indicate that the enzyme(s) that covalently attach Lpp to the cell wall may be limiting (Magnet *et al.*, 2007). Does the ratio of the two topological forms vary under any growth conditions? Can the transmembrane topological form be converted into the periplasmic form? The answers to these questions may provide insights into the physiological role of the transmembrane form of Lpp.

Certain other bacterial membrane proteins have dual properties. For example, secreted bacterial protein toxins, such as colicins and haemolysins, can adopt both soluble and membrane-inserted topologies (Parker and Feil, 2005). Proteins with dual membrane orientations, such as EmrE, also exist (Rapp *et al.*, 2007). Finally, integral OM proteins, such as OmpA and OprF, can exist in either 'open' or 'closed' barrel conformations (Sugawara and Nikaïdo, 1994; Sugawara *et al.*, 2006). By existing in two distinct subcellular locations, our work adds yet another example of a bacterial membrane protein that has dual properties. Lpp has been a useful model lipoprotein for decades and there is clearly more to be learned from this simple, 58 amino acid lipoprotein.

## Experimental procedures

### *Bacterial strains and cell culture conditions*

Bacterial cells were grown in Luria–Bertani (LB) broth or on LB agar at 37°C unless specified otherwise. Bacterial strains, most of which are derivatives of *E. coli* MC4100 (Silhavy *et al.*, 1984), are listed in Table S1.

### *Whole cell surface labelling*

Cells were subcultured 1:1000 from overnight-grown cultures into 5 ml cultures with 0.02% (w/v) arabinose, when necessary to induce expression from plasmid constructs. Cultures were grown with aeration for 3 h, after which time the cells were pelleted, washed three times in PBS, and normalized to OD<sub>600</sub> = 10.0 in PBS. NHS-LC-LC-biotin (Pierce, Cat. No. 21343) was added to a final concentration of 2% (v/v) from a 25 mg ml<sup>-1</sup> stock in dimethylsulphoxide (DMSO) and the cells were held for 20 min at room temperature. PEG-NHS-biotin was added to a final approximate concentration of 10% (v/v) from a 5% (w/v) stock in 50% DMSO and reacted for 20 min at room temperature. The reactions were quenched by the addition of Tris, pH = 7.5 to a final concentration of 250 mM. Finally, the labelled cells were washed in PBS and resuspended directly in the original reaction volume of SDS/PAGE loading buffer.

### *Subcellular fractionation*

To separate the soluble and insoluble subcellular fractions, a 100 ml mid-exponential culture was washed, normalized to OD<sub>600</sub> = 10.0 in PBS, and labelled as above. The cells were then lysed by passage through a French pressure cell at 20 000 PSI. The lysates were spun in a microcentrifuge for 1 min to remove intact cells and 1 ml of the supernatant was spun for 30 min at 100 000 r.p.m. in a Beckman TLA 100.2 ultracentrifuge rotor. The supernatant was saved as the soluble fraction and the pellet was washed with water and saved as the insoluble fraction by solubilizing the pellet in 1 ml of 1% SDS. In order to examine whether NHS-LC-LC-biotin partitions into membranes, the same process was followed,

except that cells were first washed, normalized, and mechanically lysed, and subsequently labelled and fractionated by ultracentrifugation.

### *Tests of the effects of NHS-LC-LC-biotin labelling on OM integrity*

In the following experiments, a single mid-exponential culture of cells was washed in PBS, normalized to OD<sub>600</sub> = 10.0, and split into two identical aliquots. One aliquot was labelled with NHS-LC-LC-biotin as described above, whereas the other aliquot was left untreated.

To examine the effects on viability, cells were diluted through 6, 10-fold serial dilutions in LB and approximately 1 µl of each dilution was spotted onto triplicate LB agar plates or LB agar plates containing 0.5% SDS. The highest dilution at which single colonies appeared was recorded and used to calculate the efficiency of plating. No difference was observed between the two treatments.

To examine the effects of NHS-LC-LC-biotin labelling on OM permeability, cells were plated on triplicate LB agar plates containing 0.5% SDS, and a filter disc containing 2, 5 or 10 µl of 0.5 M EDTA was placed in the centre of the plate. The plates were incubated overnight at 37°C and the zones of clearing were recorded. No difference in the size of the zone of clearing was observed between the labelled and unlabelled trials at any EDTA concentration.

To examine the effects of NHS-LC-LC-biotin labelling on OM permeability, the proportion of cells that were permeable was determined using Sytox Green at a final concentration of 5 mM according to the manufacturer's instructions (Invitrogen). Over 2000 cells for each condition were examined using light and fluorescence microscopy, and the proportion of fluorescent, permeable cells was calculated.

To examine the effects of NHS-LC-LC-biotin labelling on growth rate, aliquots of labelled and unlabelled cells were diluted 100-fold into triplicate broth cultures and the optical density was monitored over several hours. The averaged treated and untreated cells grew with indistinguishable kinetics.

### *Fractionation of the free and bound forms of Lpp*

To separate the free- and bound-form Lpp fractions, 1 ml of cells was labelled, washed, resuspended in PBS + 1% SDS, and boiled for 20 min. The intact peptidoglycan sacculi were then pelleted by spinning for 30 min at 100 000 r.p.m. in a Beckman TLA 100.2 ultracentrifuge rotor. Following this, the supernatant fraction was saved as the free-form fraction. The peptidoglycan pellets were gently washed 4 times with 10 mM Tris, pH = 7.5. Next, the pellets were resuspended in 1 ml of 10 mM Tris, pH = 7.5 and dispersed using a brief pulse from a probe tip sonicator. Following this, the sacculi were incubated overnight at 37°C in the absence or presence of 0.2 mg ml<sup>-1</sup> lysozyme to yield bound-form fraction samples. No proteins were ever observed in bound-form samples that were not treated with lysozyme, indicating that the peptidoglycan sacculi were intact and were not contaminated with free-form fraction proteins.

To examine whether bound-form Lpp could be labelled by NHS-LC-LC-biotin *in vivo*, mid-exponential cultures of CC01

and CC14 were washed and normalized as above and NHS-LC-LC-biotin was added to a final concentration of 5% (v/v). Cells were reacted, quenched and fractionated into the free- and bound-form fractions, as above. Following fractionation, blots were detected with streptavidin-HRP at 1:10 000 and exposed to film for an extended length of time.

#### Immunoblot analyses

$\alpha$ -Lpp and  $\alpha$ -biotin immunoblot analyses were carried out with standard protocols with nitrocellulose membranes in TBS-T. Prior to detection, blots were blocked for 1 h with 5% milk in TBS-T. Lpp was detected with polyclonal rabbit primary  $\alpha$ -Lpp antiserum (kindly provided by M. Inouye) at 1:400 000 and a donkey,  $\alpha$ -rabbit, HRP-conjugated secondary antibody at 1:8000 (GE Healthcare, Cat. No. NA934V). Biotin and LC-LC-biotin were detected with a streptavidin-HRP conjugate (Rockland Immunochemicals, Cat. No. S000-03) in TBS-T + 1% milk used at 1:20 000 from a 1 mg ml<sup>-1</sup> stock. OmpA was detected with an  $\alpha$ -LamB polyclonal rabbit primary antiserum that cross-reacts with OmpA at 1:20 000 in TBS-T. NlpB was detected with  $\alpha$ -NlpB polyclonal rabbit primary antiserum at 1:20 000 in TBS-T (Wu *et al.*, 2005). TraT was detected with  $\alpha$ -TraT polyclonal rabbit primary antiserum at 1:20 000 in TBS-T (see below).

#### Synthesis of PEG-NHS-biotin

PEG 3500-NHS-biotin was synthesized using tricarballylic acid, *O*-benzylhydroxylamine hydrochloride, PEG 3500, and biotin (Acros) using both previously published (Schlemminger *et al.*, 2003) and standard chemical procedures. Briefly, tricarballylic acid was condensed onto *O*-benzylhydroxylamine to yield a benzyl-protected NHS ester derivative. The free carboxylic acid of this product was esterified with PEG 3500. The product was then deprotected, activated, and biotin was linked to the activated NHS ester. A more detailed description of this synthesis is shown in Fig. S2.

#### Proteolysis methods

The susceptibility of the LC-LC-biotin label attached to Lpp and OmpA was examined by treating labelled and washed whole cells with 0.5 mg ml<sup>-1</sup> trypsin or proteinase K at 37°C for 1 h in 100 mM NaCl, 20 mM Tris, pH = 7.5. The proteases were inactivated by adding PMSF to 5 mM and washing the cells four times in 100 mM NaCl, 20 mM Tris, pH = 7.5 + 1 mM PMSF. The cells were then normalized to an equivalent OD<sub>600</sub> = 5.0 by resuspending the cell pellet in SDS/PAGE loading buffer + 1 mM PMSF.

To examine the protease susceptibility of Lpp-FLAG, *lpp* null cells (strain CC03) harbouring pBAD/*lpp* or pBAD/*lpp*-FLAG plasmids were grown to mid-log phase in LB broth supplemented with 0.02% arabinose to induce expression of the Lpp constructs. Cells were washed in 100 mM NaCl, 20 mM Tris, pH = 7.5, normalized to OD<sub>600</sub> = 10.0, and incubated in the presence of 0.5 mg ml<sup>-1</sup> trypsin + 2% glycerol at 37°C for 1 h. The cells were then washed three times in PBS, normalized to OD<sub>600</sub> = 10.0 in PBS, and labelled with NHS-LC-LC-biotin, as described above.

To examine the susceptibility of free- and bound-form Lpp to periplasmic proteolysis, cells were washed in 100 mM NaCl, 20 mM Tris, pH = 7.5 and normalized to OD<sub>600</sub> = 10.0. EDTA was added to a final concentration of 5 mM and the cells were split into two identical aliquots. Proteinase K was added to one aliquot to a final concentration of 0.5 mg ml<sup>-1</sup> and the cells were incubated at 37°C for 1 h. Following this, PMSF was added to a final concentration of 5 mM and the cells were separated into free- and bound-form fractions, as above, except that 1 mM PMSF was added to the overnight lysozyme digestion reactions.

#### Purification of TraT and production of $\alpha$ -TraT antibodies

To purify TraT, *traT* was amplified using oligonucleotides TraTNSolFor and TraTSalSTOPRev, digested with NdeI and SalI, and cloned into similarly digested pET28a. The resulting plasmid, pET/TraTNSol, was inserted into BL21( $\lambda$ DE3) cells. An overnight culture of the expression strain was subcultured 1:150 into 6 l of LB + 20  $\mu$ M IPTG and the cells were grown for 3 h with shaking. Cells were pelleted, washed and mechanically lysed using a pressure lysis device (Microfluidics). His-TraT was batch-purified from the cells using 12 ml Ni-NTA resin, according to the manufacturer (Qiagen). The 100 and 250 mM imidazole eluate fractions were pooled and dialysed against two large volumes of 100 mM NaCl, 50 mM Tris, 0.5 mM EDTA, 5% glycerol, pH = 7.75. Finally, the dialysed protein was concentrated by passage through a 50 kDa MWCO Centricon filter (Millipore) and retention on a 30 kDa MWCO Centricon filter.

One milligram purified His-TraT was mixed with Freund's complete adjuvant and injected into a New Zealand white rabbit (Covance). Serum was collected at 28 days and reacted with a 25 kDa protein in F+ cells, but showed no cross-reactivity with other *E. coli* proteins in F- cells.

#### Molecular biological methods

The oligonucleotides used in this study are listed in Table S1. Chromosomal mutant *lpp* alleles were constructed in CC05 using single oligo-mediated mutagenesis (Costantino and Court, 2003) with oligos 'delta17-37ultramer' and 'lppdeltaK58.' CC06 (an *lpp* <sub>$\Delta$ 17-37</sub> strain) was identified by screening candidate colonies with PCR reactions using 'lppF2' and 'lppseqrev' primers to identify strains with PCR amplicons that were shorter than the wild-type *lpp* product. CC07 (an *lpp* <sub>$\Delta$ K58</sub> strain) was identified by screening candidate colonies for sensitivity to 0.5% SDS + 1 mM EDTA. The sequence of both alleles was determined by PCR amplifying the *lpp* gene using primers 'lppF2' and 'lppseqrev' and then sequencing the product with primer *lpp* 'lppF3' at GENEWIZ (South Plainfield, NJ, USA) to verify the expected sequence from the gene's promoter to the stop codon. Following this, the alleles were mobilized from CC06 and CC07 into CC01 by P1 transduction to create strains CC08 and CC09 (*lpp* <sub>$\Delta$ 17-37</sub> and *lpp* <sub>$\Delta$ K58</sub> strains respectively).

Standard molecular biological methods were used for plasmid construction. Plasmids pBAD/*lpp*, pBAD/*lpp* <sub>$\Delta$ K58</sub> and pBAD/*lpp* <sub>$\Delta$ 55-58</sub> were constructed by amplifying the *lpp* open reading frame and ribosome binding site using primers

pBADlppFor and either pBADlppRev, pBADlpp-CKRev or pBADlpp-C2K rev respectively. The correct sequence of all plasmids was determined by directly sequencing with both pBAD-Forward and pBAD-Reverse primers at GENEWIZ.

#### Examination of NHS-LC-LC-biotin surface specificity using NlpB and TraT control lipoproteins

To examine the surface specificity of NHS-LC-LC-biotin using lipoproteins with known orientations, CC11 (*ompA*-null, *nlpB*-null) cells were transformed with pBAD/nlpB or pBAD/traT. An *ompA*-null strain was used, as NlpB co-migrates with OmpA. NlpB expression was induced with 0.02% arabinose and TraT expression was induced with 0.001% or 0.002% arabinose, and the cells were surface-labelled with NHS-LC-LC-biotin, as described above. Known quantities of purified NlpB-His (kindly provided by J. Malinverni) or His-TraT were used as references for immunoblot quantification of each protein's expression level in the labelled samples. Reference protein standards were varied through several, twofold dilutions, and were used in ranges that flanked the concentration of the experimental proteins. Duplicate quantifications were performed for each sample.

#### Mass spectrometric analyses

For mass spectrometric analyses, whole cells were surface-labelled with NHS-LC-LC-biotin and insoluble-fraction SDS/PAGE samples were prepared as described above. The samples were separated using 10–20% SDS/PAGE gels and the 10–12 kDa region of the gel was excised and diced into 1-mm<sup>3</sup> pieces. The sample was digested with 12.5 ng ml<sup>-1</sup> sequencing grade modified trypsin (Promega), and the resulting peptides were extracted on reverse-phase resin (Poros 20 R2, PerSeptive Biosystems), as previously described (Cristea *et al.*, 2005). The peptide mixtures were eluted directly onto a MALDI target in 50% methanol, 20% acetonitrile, 0.1% trifluoroacetic acid + 2 mg ml<sup>-1</sup>  $\alpha$ -Cyano-4-hydroxycinnamic acid matrix. MALDI MS and MS/MS analyses were performed using a MALDI LTQ Orbitrap XL, as previously described (Luo *et al.*, 2010). Lists of the expected m/z for tryptic peptides and ion fragments were generated using Prowl (<http://prowl.rockefeller.edu/>) with K + 452.24571 for modification by LC-LC-biotin.

For MALDI LTQ MS<sup>2</sup> analysis, the ion of interest (LDNMTK\*YRK\*, m/z 2144.143) was isolated in the ion trap with a 2 Da mass range window, and subjected to collision induced dissociation (CID), as described (Luo *et al.*, 2010), using a normalized collision energy of 35%. As expected from the primary sequence of this ion, the predominant ion was the y<sub>8</sub> ion, resulting from the preferential cleavage at the carboxyl-terminus of the aspartic acid (Qin and Chait, 1995). Thus, this ion (NMTK\*YRK\*, m/z 1916.11) was further subjected to MALDI LTQ CID MS<sup>3</sup> analysis with the same settings as above (Fig. S1).

#### In vivo cross-linking

To cross-link Lpp *in vivo*, cells were subcultured 1:1000 and grown for 3 h, washed three times in PBS, and normalized to

OD<sub>600</sub> = 10.0. Whole cells were cross-linked with a range of concentrations of disuccinimidyl suberate concentrations for 20 min at room temperature. Following this, the reactions were quenched by the addition of Tris, pH = 7.5 to a final concentration of 250 mM. Finally, the cross-linked cells were washed in PBS and resuspended in SDS/PAGE loading buffer at their original concentration.

#### Acknowledgements

This work was supported by an NIH postdoctoral fellowship AI07402 (C.E.C.) and NIH grants GM34821 (T.J.S.) and DP1DA026192 (I.M.C.). We give our most sincere thanks to members of the Silhavy lab, R. Young, Z. Gitai, J. Werner and K. Cowles for helpful discussions and careful reading of the manuscript, to J. Malinverni for pBAD/nlpB and purified NlpB protein, to V. Carabetta for assistance with MALDI sample preparation, C. Shebelut and K. Cowles for assistance with microscopy, and to L. Shi for media preparation.

#### References

- Behr, M.G., Schnaitman, C.A., and Pugsley, A.P. (1980) Major heat-modifiable outer membrane protein in Gram-negative bacteria: comparison with the OmpA protein of *Escherichia coli*. *J Bacteriol* **143**: 906–913.
- Bolla, J.M., Bernadac, A., Lazdunski, C., and Pages, J.M. (1990) A genetic engineering approach to study the mode of assembly of the OmpF porin in the envelope of *E. coli*. *Biochimie* **72**: 385–395.
- Braun, V., and Bosch, V. (1972) Sequence of the murein-lipoprotein and the attachment site of the lipid. *Eur J Biochem* **28**: 51–69.
- Braun, V., and Rehn, K. (1969) Chemical characterization, spatial distribution, and function of a lipoprotein (murein-lipoprotein) of the *E. coli* cell wall. The specific effect of trypsin on the membrane structure. *Eur J Biochem* **10**: 426–438.
- Braun, V., and Sieglin, U. (1970) The covalent murein-lipoprotein structure of the *Escherichia coli* cell wall. The attachment site of the lipoprotein on the murein. *Eur J Biochem* **13**: 336–346.
- Braun, M., and Silhavy, T.J. (2002) Imp/OstA is required for cell envelope biogenesis in *Escherichia coli*. *Mol Microbiol* **45**: 1289–1302.
- Braun, V., and Wolff, H. (1970) The murein-lipoprotein linkage in the cell wall of *Escherichia coli*. *Eur J Biochem* **14**: 387–391.
- Chen, R., Schmidmayr, W., Kramer, C., Chen-Schmeisser, U., and Henning, U. (1980) Primary structure of the major outer membrane protein II\* (*ompA* protein) of *Escherichia coli*. *Proc Natl Acad Sci USA* **77**: 4592–4596.
- Choi, D.-S., Yamada, H., Mizuno, T., and Mizushima, S. (1986) Trimeric structure and localization of the major lipoprotein in the cell surface of *Escherichia coli*. *J Biol Chem* **261**: 8953–8957.
- Cole, S.T., Chen-Schmeisser, U., Hindennach, I., and Henning, U. (1983) Apparent bacteriophage-binding region of an *Escherichia coli* K-12 outer membrane protein. *J Bacteriol* **153**: 581–587.

- Cornelissen, C.N., Kelley, M., Hobbs, M.M., Anderson, J.E., Cannon, J.G., Cohen, M.S., and Sparling, P.F. (1998) The transferrin receptor expressed by gonococcal strain FA1090 is required for the experimental infection of human male volunteers. *Mol Microbiol* **27**: 611–616.
- Costatino, N., and Court, D.T. (2003) Enhanced levels of lambda Red-mediated recombinants in mismatch repair mutants. *Proc Natl Acad Sci USA* **100**: 15748–15753.
- Cristea, I.M., Williams, R., Chait, B.T., and Rout, M.P. (2005) Fluorescent proteins as proteomic probes. *Mol Cell Proteomics* **4**: 1933–1941.
- Dashper, S.G., Hendtlass, A., Slakeski, N., Jackson, C., Cross, K.J., Brownfield, L., et al. (2000) Characterization of a Novel Outer Membrane Hemin-Binding Protein of *Porphyromonas gingivalis*. *J Bacteriol* **182**: 6456–6462.
- Datta, D.B., Arden, B., and Henning, U. (1977) Major proteins of the *Escherichia coli* outer cell envelope membrane as bacteriophage receptors. *J Bacteriol* **131**: 821–829.
- Decad, G.M., and Nikaido, H. (1976) Outer membrane of Gram-negative bacteria XII. Molecular-sieving function of cell wall. *J Bacteriol* **128**: 325–336.
- Drummelsmith, J., and Whitfield, C. (2000) Translocation of group 1 capsular polysaccharide to the surface of *Escherichia coli* requires a multimeric complex in the outer membrane. *EMBO J* **19**: 57–66.
- Hirota, Y., Suzuki, H., Nishimura, Y., and Yasuda, S. (1977) On the process of cellular division in *Escherichia coli*: a mutant of *E. coli* lacking a murein-lipoprotein. *Proc Natl Acad Sci USA* **74**: 1417–1420.
- Inouye, M. (1974) A three-dimensional molecular assembly model of a lipoprotein from the *Escherichia coli* outer membrane. *Proc Natl Acad Sci USA* **71**: 2396–2400.
- Inouye, M., Shaw, J., and Shen, C. (1972) The assembly of a structural lipoprotein in the envelope of *Escherichia coli*. *J Biol Chem* **247**: 8154–8159.
- Jin, S., Joe, A., Lynett, J., Hani, E.K., Sherman, P., and Chan, V.L. (2001) JlpA, a novel surface-exposed lipoprotein specific to *Campylobacter jejuni*, mediates adherence to host epithelial cells. *Mol Microbiol* **39**: 1225–1236.
- Leuzzi, R., Serino, L., Scarselli, M., Savino, S., Fontana, M., Monaci, E., et al. (2005) Ng-MIP, a surface-exposed lipoprotein of *Neisseria gonorrhoeae*, has peptidyl-prolyl cis/trans isomerase (PPIase) activity and is involved in persistence in macrophages. *Mol Microbiol* **58**: 669–681.
- Li, S.J., and Cronan, J.E. (1992) The gene encoding the biotin carboxylase subunit of *Escherichia coli* acetyl-CoA carboxylase. *J Biol Chem* **267**: 855–863.
- Luo, Y., Li, T., Yu, F., Kramer, T., and Cristea, I.M. (2010) The performance of a MALDI LTQ Orbitrap and its application to the study of protein complexes. *J Am Soc Mass Spectrom* **21**: 34–36.
- Magnet, S., Bellais, S., Dubost, L., Fourgeaud, M., Mainardi, J.-L., Petit-Frere, S., et al. (2007) Identification of the L,D-transpeptidases responsible for attachment of the braun lipoprotein to *Escherichia coli* peptidoglycan. *J Bacteriol* **189**: 3927–3931.
- Manning, P.A., Beutin, L., and Achtman, M. (1980) Outer membrane of *Escherichia coli*: properties of the F sex factor *traT* protein which is involved in surface exclusion. *J Bacteriol* **142**: 285–294.
- Nakae, T., and Nikaido, H. (1975) Outer Membrane as a Diffusion Barrier in *Salmonella typhimurium*. Penetration of oligo- and polysaccharides into isolated outer membrane vesicles and cells with degraded peptidoglycan layer. *J Biol Chem* **250**: 7359–7365.
- Neidhardt, F.C. (1996) Chemical composition of *Escherichia coli*. In *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Neidhardt, F.C., Curtis, R.I., Ingraham, J.L., Lin, E.C.C., Low, K.B., Magasnik, B., et al. (eds). Washington, DC: ASM Press, pp. 13–16.
- Nikaido, H. (1996) Outer membrane. In *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Neidhardt, F.C., Curtis, R.I., Ingraham, J.L., Lin, E.C.C., Low, K.B., Magasnik, B., et al. (eds). Washington, DC: ASM Press, pp. 29–47.
- Nikaido, H. (2003) Molecular basis of bacterial outer membrane permeability revisited. *Microbiol Mol Biol Rev* **67**: 593–656.
- Nikaido, H., and Nakae, T. (1979) The outer membrane of Gram-negative bacteria. *Adv Microb Physiol* **20**: 163–250.
- Nikaido, H., and Rosenberg, E.Y. (1981) Effect of solute size on diffusion rates through the transmembrane pores of the outer membrane of *Escherichia coli*. *J Gen Physiol* **77**: 121–135.
- Parker, M.W., and Feil, S.C. (2005) Pore-forming protein toxins: from structure to function. *Prog Biophys Mol Biol* **88**: 91–142.
- Qin, J., and Chait, B.T. (1995) Preferential fragmentation of protonated gas-phase peptide ions adjacent to acidic amino acid residues. *J Am Chem Soc* **117**: 5411–5412.
- Rapp, M., Seppala, S., Granseth, E., and von Heijne, G. (2007) Emulating membrane protein evolution by rational design. *Science* **315**: 1282–1284.
- Robinson, L.S., Ashman, E.M., Hultgren, S.J., and Chapman, M.R. (2006) Secretion of curli fibre subunits is mediated by the outer membrane-localized CsgG protein. *Mol Microbiol* **59**: 870–881.
- Ronco, J., Charbit, A., and Hofnung, M. (1988) Creation of targets for proteolytic cleavage in the LamB protein of *E. coli* K12 by genetic insertion of foreign sequences. *Biochimie* **72**: 183–189.
- Ruiz, N., Kahne, D., and Silhavy, T.J. (2006) Advances in understanding bacterial outer-membrane biogenesis. *Nat Rev Microbiol* **4**: 57–66.
- Schlemminger, I., Mole, D.R., McNeill, L.A., Dhanda, A., Hewitson, K.S., Tian, Y.-M., et al. (2003) Analogues of dealanylalohopcin are inhibitors of human HIF prolyl hydroxylases. *Bioorg Med Chem Lett* **13**: 1451–1454.
- Schulze, R.J., and Zuckert, W.R. (2006) *Borrelia burgdorferi* lipoproteins are secreted to the outer surface by default. *Mol Microbiol* **59**: 1473–1484.
- Shu, W., Liu, J., Ji, H., and Lu, M. (2000) Core structure of the outer membrane lipoprotein from *Escherichia coli* at 1.9 angstrom resolution. *J Mol Biol* **299**: 1101–1112.
- Silhavy, T.J., Berman, M.L., and Enquist, L.W. (1984) *Experiments with Gene Fusions*. Plainview, NY: Cold Spring Harbor Lab Press.
- Sugawara, E., and Nikaido, H. (1994) OmpA protein of *Escherichia coli* outer membrane occurs in open and closed channel forms. *J Biol Chem* **269**: 17981–17987.

- Sugawara, E., Nestorovich, E.M., Bezrukov, S.M., and Nikaido, H. (2006) *Pseudomonas aeruginosa* porin OprF exists in two different conformations. *J Biol Chem* **281**: 16220–16229.
- Suzuki, H., Nishimura, Y., Yasuda, S., Nishimura, A., Yamada, M., and Hirota, Y. (1978) Murein-lipoprotein of *Escherichia coli*: a protein involved in the stabilization of bacterial cell envelope. *Mol Gen Genet* **167**: 1–9.
- Tokuda, H. (2009) Biogenesis of outer membranes in Gram-negative bacteria. *Biosci Biotechnol Biochem* **73**: 465–473.
- Wu, T., Malinverni, J., Ruiz, N., Kim, S., Silhavy, T.J., and Kahne, D. (2005) Identification of a multicomponent complex required for outer membrane biogenesis in *Escherichia coli*. *Cell* **121**: 235–245.
- Yem, D.E., and Wu, H.C. (1977) Physiological characterization of an *Escherichia coli* mutant altered in the structure of murein lipoprotein. *J Bacteriol* **133**: 1419–1426.
- Yokota, N., Kuroda, T., Matsuyama, S., and Tokuda, H. (1999) Characterization of the LolA-LolB system as the general lipoprotein localization mechanism of *Escherichia coli*. *J Biol Chem* **274**: 30995–30999.
- Zhang, W.-Y., and Wu, H.C. (1992) Alterations of the carboxyl-terminal amino acid residues of *Escherichia coli* lipoprotein affect the formation of murein-bound lipoprotein. *J Biol Chem* **267**: 19560–19564.
- Zhang, W.-Y., Dai, R.-M., and Wu, H.C. (1992) Deletion of internal twenty-one amino acid residues of *Escherichia coli* prolipoprotein does not affect the formation of the murein-bound lipoprotein. *FEBS Lett* **311**: 311–314.

### Supporting information

Additional supporting information may be found in the online version of this article.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.